# **UMBC Intercomparisons**

# Ground-based Lidar, BAM and Nephelometer vs. Aircraft HSRL, in-situ SO<sub>4</sub><sup>2-</sup> and f(RH)

Patricia Sawamura, Raymond M. Hoff. Ruben Delgado, Timothy Berkoff, Bruce Anderson, Richard Ferrare, Chris Hostetler, John Hair, Ray Rogers, Mike Obland




# $P(\lambda_0,z) = \frac{KO(\lambda_0,z)}{z^2} \beta(\lambda_0,z) \exp[-2\int\limits_0^z \alpha(\lambda_0,z')dz']$ Overall, extinction retrievals from ELF agree very well with HSRL measurements. Cases in which good agreement


occurs indicate very well mixed aerosols within the PBL,

when S does not vary with height.

 $P(\lambda_{Ra}, z) = \frac{K'O(\lambda_{Ra}, z)}{z^2} \beta(\lambda_{Ra}, z) \exp\left[-\int_0^z (\alpha(\lambda_0, z') + \alpha(\lambda_{Ra}, z'))dz'\right]$   $\alpha_{aer}(\lambda_0, z) = \frac{\frac{d}{dz} \ln\left[\frac{N_{Ra}(z)}{P(\lambda_{Ra}, z)z^2}\right] - \alpha_{mol}(\lambda_0, z)}{1 + \left(\frac{\lambda_0}{\lambda_{Ra}}\right)^{a(z)}}$ 


$$\begin{split} P(\lambda_0,z) &= \frac{KO(\lambda_0,z)}{z^2} \, \beta(\lambda_0,z) \exp[-2\int\limits_0^z S(\lambda_0,z') \beta(\lambda_0,z') dz'] \\ &= \underbrace{\text{Extinction-to-Backscatter}}_{\text{or Lidar Ratio (S)}} S(\lambda_0,z) \cong S(\lambda_0) = \frac{\alpha_{aer}(\lambda_0,z)}{\beta_{aer}(\lambda_0,z)} \end{split}$$





### Assumptions:


- -- BAM and NEPH measure dry aerosols;
- -- ELF measures aerosols at ambient conditions;



## Specific Scattering Coefficient (SSC):

$$SSC = \frac{Scattering}{MassConc.} = 4.8 \pm 0.4 [m^2 g^{-1}]$$

Extinction was dominated by scattering Neph = 1.23(12)xELF + 41(13), R<sup>2</sup> = 0.63



$$\alpha_{dry} = \frac{\alpha_{amb}}{f(RH)} = SSC \times PM \ 2.5$$

$$f(RH)_{UMBC} = \frac{6.5}{4.8} = 1.35$$

$$f(RH)_{P3B} = 1.7 \pm 0.2 \approx f(RH)_{UMBC} - 1.7\sigma_{P3B}$$

### P3B UMBC Overpasses at 372m ± 31m

### Acknowledgements:

NASA cooperative agreements: NNX10AR38G (DISCOVER-AQ) and NNX10AT36A(JCET Task 336)